

University of Groningen

A Simple Road for the Transformation of Few-Layer Graphene into MWNTs

Quintana, Mildred; Grzelczak, Marek; Spyrou, Konstantinos; Calvaresi, Matteo; Bals, Sara; Kooi, Bart; Van Tendeloo, Gustaaf; Rudolf, Petra; Zerbetto, Francesco; Prato, Maurizio

Published in:
Journal of the American Chemical Society

DOI:
[10.1021/ja303131j](https://doi.org/10.1021/ja303131j)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Quintana, M., Grzelczak, M., Spyrou, K., Calvaresi, M., Bals, S., Kooi, B., Van Tendeloo, G., Rudolf, P., Zerbetto, F., & Prato, M. (2012). A Simple Road for the Transformation of Few-Layer Graphene into MWNTs. *Journal of the American Chemical Society*, 134(32), 13310-13315.
<https://doi.org/10.1021/ja303131j>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supporting Information

for

A Simple road for the transformation of few-layer graphene into MWNTs




Mildred Quintana,¹ Marek Grzelczak,¹ Konstantinos Spyrou,² Matteo Calvaresi,³ Sara Bals,⁴ Bart Kooi,² Gustaaf van Tendeloo,⁴ Petra Rudolf,² Francesco Zerbetto³ and Maurizio Prato¹

1. Center of Excellence for Nanostructured Materials (CENMAT) and INSTM, unit of Trieste,
Dipartimento di Scienze Chimiche e Farmaceutiche, University of Trieste, Piazzale Europa 1, I-
34127 Trieste, Italy


2. Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, NL-9747AG
Groningen, The Netherlands

3. Dipartimento di Chimica “G. Ciamician”, Università di Bologna, V. F. Selmi 2, 40126 Bologna,
Italy


4. EMAT, University of Antwerp, Groenenborgerlaan 171, B-2020 ANTWERP, Belgium




SI-1. MWNT formation on the edges of graphene layers. Evolution during the time is shown.




SI-2. HR-TEM of graphene edges and MWNTs formation.



SI-3. MWNTs produced by the addition on **Fc-CHO** mixed with graphene layers.





SI-4. G-Fc-CHO





SI-5. O 1s peak of D-Fc-CHO



Control experiments preparation. **G-1** dispersion was used as starting material for the control experiments. As control experiment, we performed a 3 h sonication of **G-1** adding (i) Fc, (ii) benzaldehyde, (iii) Fc together with benzaldehyde, (iv) Fc-COOH, and (v) FcCON(CH₃)₂. The product was copiously washed by filtration with fresh DMF in order to remove antioxidant and by-product molecules. Samples were re-dispersed in a bath ultrasonicator (few seconds) in 10 ml of fresh DMF. Centrifugation of all dispersions was carried out at 3000 rpm during 30 min. Two liquid fractions, of 5 ml each, of the **D-Fc-CHO** dispersion were collected and analyzed.





SI-6. Control experiment Fc




SI-7- Control experiment by adding benzaldehyde.



SI-8. Control experiment by adding Fc and benzaldehyde.



SI-9. Control experiment by adding Fc-COOH



SI-10. Control experiment by adding $\text{Fc-CON}(\text{CH}_3)_2$.

Molecular Mechanics and Molecular Dynamics calculations for an increasing number of Fc-CHO molecules deposited on the nanoribbon obtained by unzipping a (8,8) CNT


To locate the minimum energy path (MEP) for the rolling up of the graphene nanoribbon in the absence or in the presence of a diverse number of Fc-CHO molecules (Fig SI-11) we carried out different molecular dynamics simulations constraining the distance between the two edges of the nanoribbon.

To avoid that the systems remain trapped in local minima, we employed a high temperature quenched molecular dynamics protocol where the system is first heated from 0 to 600K in 100 ps. Then, a trajectory of 10 ns is carried out at constant temperature (600 K). The RATTLE algorithm constrained the stretching of bonds with hydrogen atoms and allowed for a timestep of 2 fs. The coordinates of the structures were saved every 10 ps for a total of 1000 conformations that were used in further minimizations. Each structure was energy minimized until the root mean square of the gradient was less than $0.01 \text{ kcal mol}^{-1}$.


To describe the structure and energetics of the system, we employed Brenner potential for nanoribbon/nanotube and MM3 potential for Fc-CHO and graphene/NT-Fc-CHO interactions, a well tested protocol using a modified version of Tinker.¹

Table SI-1. Energy barrier for a SWNT (8,8) rolling up with different number of Fc-CHO molecules.


N° of Fc-CHO molecules	Energy barrier for a SWNT (8,8) rolling	Stabilization energy of the Fc-CHO molecule (Kcal mol ⁻¹)
0	120.3	
1	102.5	17.8
2	86.0	16.5
3	72.0	14.0
4	63.3	8.7



SI-11. Minimum Energy Path (MEP) for the rolling up of the graphene nanoribbon in the absence (purple line) and in the presence of four Fc-CHO molecules (yellow line).



SI-12. Representative snapshots during rolling up of the graphene nanoribbon in the absence and in the presence of four Fc-CHO molecules.



SI-13. MWNTs after annealing treatment.


Calculation of the Electron affinities (Eas)

The computations reported in table SI-2 were performed with the Gaussian09² series of programs. The geometry of the various critical points was fully optimized with the gradient method available in Gaussian 09 at the DFT level using the M06 functional³ and the 6-31g** basis set.⁴ Single point calculation were then carried out using AUG-cc-PVTZ basis set on the previously optimized geometries.⁵ Solvent effect (DMF, $\epsilon = 37.219$) was taken in account during optimization and single point calculations using the Polarizable Continuum Model (PCM) in the integral equation formalism variant (IEFPCM).⁶ The solute cavity was built up using the UFF radii,⁷ which place a sphere around each solute atom, with the radii scaled by a factor of 1.1.

The electron affinities (Eas) were calculated as the energy difference between anionic optimised structure and neutral optimised one.

Table SI-2. Electron affinities (Eas).

Molecule	EA (kcal/mol)
Fc-CHO	-50.0
Fc-COOH	-44.4
Fc-CON(CH ₃) ₂	-32.0
Fc	-31.5
Benzaldehyde	-26.6



SI-14. Isosurfaces spin densities of Fc-CHO (left) and Fc (right) radical molecules plotted using an isovalue of $0.004 \text{ } ea_0^{-3}$.

- 1) Plank, W.; Pfeiffer, R.; Schaman, C.; Kuzmany, H.; Calvaresi, M.; Zerbetto, F.; Meyer, J. *ACS Nano*, **2010**, *4*, 4515–4522.
- 2) Gaussian 09, Revision **A.02**, Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, N. J.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian, Inc., Wallingford CT, 2009.
- 3) Zhao, Y.; Truhlar, D. *Theor. Chem. Acc.*, **2008**, *120*, 215-241.
- 4) Frisch, M. J.; Pople, J. A.; Binkley, J. S. *J. Chem. Phys.*, **1984**, *80*, 3265–3269.
- 5) Dunning Jr., T. H. *J. Chem. Phys.*, **1989**, *90*, 1007-1023.
- 6) Tomasi, J.; Mennucci, B.; Cammi, R. *Chem. Rev.*, **2005**, *105*, 2999–3093.
- 7) Rappè, A. K.; Casewit, C. J.; Colwell, K. S.; Goddard III, W. A.; Skid, W. M. *J. Am. Chem. Soc.* **1992**, *114*, 10024–10039.